

On The Inhomogeneous Quantum Groups

A. Shariati ^{†,‡}, A. Aghamohammadi ^{◇,‡}

[†] *Department of Physics, Sharif University of Technology, P.O.Box 9161, Tehran 11365, Iran.*

[‡] *Institute for Studies in Theoretical Physics and Mathematics, P.O.Box 5746, Tehran 19395, Iran.*

[◇] *Department of Physics, Alzahra University P.O.Box 19834, Tehran, Iran*

Abstract

Using the multi-parametric deformation of the algebra of functions on $GL(n+1)$ and the universal enveloping algebra $U(\mathfrak{gl}(n+1))$, we construct the multi-parametric quantum groups $IGL_q(n)$ and $U_q(\mathfrak{gl}(n))$.

The quantization of inhomogeneous groups is not well established. These non-semisimple groups are important, both from a purely mathematical point of view, and also because one of the most important groups appearing in physics, viz. the Poincaré group, is inhomogeneous. Various authors in the last few years studied the quantization of these objects from different viewpoints and by different methods [1-7].

In this paper we consider the inhomogeneous general linear group $IGL(n)$. We will show that in analogy with the classical case, the multi-parameteric deformation of the parent $GL(n+1)$ will lead to a consistent quantization of both the algebra of functions and the universal enveloping algebra of the inhomogeneous group $IGL(n)$. Multi-parametric quantum groups have been studied by various authors [8-14]. In this paper we will use the notations of Ref. [13]. To clarify our approach, we will first consider the two dimensional case $IGL(2)$; then we will generalize to the higher dimensions. Throughout this paper, when we use a subscript q for an object, we mean the multi-parameteric deformation of that object. The number of, and the conditions on, the parameters will be clear from the context.

Let us consider the two dimensional inhomogeneous group $IGL(2)$. Its action on the two di-

mensional plane $\mathbb{R}^2$ is:

$$x \mapsto x' = ax + by + u \quad y \mapsto y' = cx + dy + v. \quad (1)$$

This action may be represented on the $z = 1$ subset of $\mathbb{R}^3$ by means of the following matrix:

$$\begin{pmatrix} x \\ y \\ 1 \end{pmatrix} \mapsto \begin{pmatrix} x' \\ y' \\ 1 \end{pmatrix} = \begin{pmatrix} a & b & u \\ c & d & v \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \\ 1 \end{pmatrix}. \quad (2)$$

From this representation we can easily write the Lie algebra spanned by:

$$\begin{aligned} H_1 &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix} & X_1^+ &= \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} & X_1^- &= \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \\ X_2^+ &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix} & X_{12}^+ &= \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} & K &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}. \end{aligned} \quad (3)$$

The set $\{H_1, X_1^+, X_1^-\}$ generates $SL(2)$; K generates dilatations (scaling), and, X_2^+ and X_{12}^+ are the generators of the translations. Note that this Lie algebra, $\mathfrak{igl}(2)$, is a subalgebra of the $\mathfrak{gl}(3)$, with the other generators of $\mathfrak{gl}(3)$ being,

$$H_2 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix} \quad X_2^- = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \quad X_{12}^- = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}. \quad (4)$$

The identity matrix I , which is a generator of $\mathfrak{gl}(3)$, is not an independent operator and we have

$$I = \frac{3}{2}K - \frac{1}{2}H_1 - H_2. \quad (5)$$

We would like to study the quantum deformed version of these relations. First of all, we have to consider a two dimensional quantum space $\mathbb{R}_q^2$ spanned by (x, y) . If this non-commutative space is to be embedded in a three dimensional quantum space $\mathbb{R}_q^3$ with coordinates (x, y, z) by setting $z = 1$, then, z must commute with everything, otherwise it would be meaningless to set $z = 1$. This is only possible if we use the multi-parameteric quantum space $\mathbb{R}_q^3$ with a special choice of the parameters. To see this we write the commutation relations of the coordinates of $\mathbb{R}_q^3$:

$$x_i x_j = q p_{ij} x_j x_i \quad i < j \quad (6)$$

Therefore, $x_3 = z$ is central only if $p_{13} = p_{23} = q^{-1}$.

The multi-parameteric quantum group $\text{GL}_q(3)$ [13] is generated by the entries of the following T matrix.

$$T = \begin{pmatrix} a & b & u \\ c & d & v \\ e & f & w \end{pmatrix}. \quad (7)$$

If we set $e = f = 0$, then w will automatically commute with a, b, c and d . However, w has non-trivial commutation relations with u and v . To make w central, we have to impose the same condition $p_{13} = p_{23} = q^{-1}$. we call the resulting quantum group, $\text{IGL}_q(2)$.

For completeness we write the commutation relations among the generators,

$$\begin{aligned} ab &= qp^{-1}ba & au &= q^2ua & bu &= q^2ub \\ cd &= qp^{-1}dc & cv &= q^2vc & dv &= q^2vd \\ ac &= qpca & bd &= qpdb & uv &= qpvu \\ bc &= p^2cb & uc &= q^{-1}pcu & ud &= q^{-1}pdu \\ ad - da &= \lambda p^{-1}bc & av - qpva &= q\lambda uc & bv - qpvb &= q\lambda ud \\ [w, \dots] &= 0 \end{aligned} \quad (8)$$

where $p := p_{12}$. The co-product, co-unity and the antipode for this quantum group are also inherited from $\text{GL}_q(3)$.

Generalizing this to n dimensions is straightforward. Use the multi-parameteric quantum space $\mathbb{R}_q^{n+1}$ and the corresponding quantum group $\text{GL}_q(n+1)$, then set $p_{in+1} = q^{-1}$ for $i = 1, 2, \dots, n$; we set $T_{n+1i} = 0$ for $i = 1, 2, \dots, n$ and then because of the choice of parameters, $x_{n+1} \in \mathbb{R}_q^{n+1}$ and $w := T_{n+1n+1} \in \text{GL}_q(n+1)$ will become central. Setting $x_{n+1} = 1$ and $w = 1$, one gets the quantum group $\text{IGL}_q(n)$.

Now we turn to the quantization of the universal enveloping algebra $U(\mathfrak{gl}(2))$. There exists a multi-parameteric deformation of $U(\mathfrak{gl}(3))$ which we name $U_q(\mathfrak{gl}(3))$. This Hopf algebra is generated by 9 generators $H_1, H_2, X_1^\pm, X_2^\pm, X_{12}^\pm$ and I . Following the analogy with the classical case, we expect that in this quantum case too, the subalgebra generated by $\{H_1, K, X_1^\pm, X_2^\pm, X_{12}^\pm\}$ is the quantum group $U_q(\mathfrak{gl}(2))$. Let's examine this idea.

The commutation relations: X_2^- and X_{12}^- do not appear in the commutation relations of the set $\{H_1, K, X_1^\pm, X_2^\pm, X_{12}^\pm\}$ where K is given by (5).

$$\begin{aligned} [H_1, K] &= 0 & [K, X_1^\pm] &= 0 \\ [K, X_2^\pm] &= 0 & [X_1^\pm, X_{12}^\pm]_q &= 0 \\ [X_2^\pm, X_{12}^\pm]_{q^{-1}} &= 0 & [H_1, X_1^\pm] &= \pm 2X_1^\pm \\ [H_1, X_{12}^\pm] &= X_{12}^\pm & [K, X_{12}^\pm] &= X_{12}^\pm \\ [X_1^\pm, X_1^\mp] &= [H_1]_q & [X_1^\pm, X_{12}^\mp] &= q^{\frac{1}{2} \pm H_1} X_2^\pm. \end{aligned} \quad (9)$$

Here we have used the notations:

$$[x]_q := \frac{q^x - q^{-x}}{q - q^{-1}} \quad (10)$$

$$[x, y]_q := q^{\frac{1}{2}} x y - q^{-\frac{1}{2}} y x. \quad (11)$$

So as an algebra $U_q(\mathfrak{igl}(2))$ is a sub-algebra of $U_q(\mathfrak{gl}(3))$ but as Hopf algebra, it isn't a sub-Hopf-algebra of $U_q(\mathfrak{gl}(3))$. Let's consider the co-product of simple roots of the multi-parametric $U_q(\mathfrak{gl}(n+1))$, [13].

$$\Delta(X_i^\pm) = X_i^\pm \otimes q^{\frac{1}{2}H_i} \left(\prod_{j=1}^n s_{ij}^{\pm \sum_{k=1}^n a_{jk}^{-1} H_k} \right) s_i^{\pm I} + q^{-\frac{1}{2}H_i} \left(\prod_{j=1}^n s_{ij}^{\mp \sum_{k=1}^n a_{jk}^{-1} H_k} \right) s_i^{\mp I} \otimes X_i^\pm \quad (12)$$

where $[a_{ij}]$ is the Cartan matrix of $\mathfrak{sl}(n+1)$, and a_{ij}^{-1} stands for ij th element of inverse of cartan matrix and their explicit form are as below

$$a_{ij} = 2\delta_{ij} - \delta_{i+1,j} - \delta_{i-1,j} \quad (13)$$

$$a_{ij}^{-1} = \frac{j(n+1-i)}{n+1} - (j-1)\theta_{ij}, \quad \theta_{ij} = \begin{cases} 1 & i < j \\ 0 & i \geq j \end{cases} \quad (14)$$

s_{ij} and s_i are extra parameters of deformation. $s_{ij} = s_{ji}^{-1}$ and $s_{ii} = 1$. Therefore, the number of parameters is $1 + \frac{1}{2}n(n+1)$. As we see the generators H_2 and I appear in the co-product of $X_1^\pm$. If we fix the parameters in a special way, then extra generators in the co-product disappear. We can construct $U_q(\mathfrak{igl}(2))$ as a sub-Hopf algebra of $U_q(\mathfrak{gl}(3))$ with the following choice for parameters.

$$s_{12} := s, \quad s_1 = s^{\frac{2}{3}}, \quad s_2 = q^{\frac{1}{2}} s^{\frac{-1}{3}} \quad (15)$$

Then the co-products of the generators of $U_q(\mathfrak{igl}(2))$ are

$$\begin{aligned} \Delta(X_1^\pm) &= X_1^\pm \otimes q^{\frac{1}{2}H_1} s^{\pm \frac{1}{2}(H_1-K)} + q^{-\frac{1}{2}H_1} s^{\mp \frac{1}{2}(H_1-K)} \otimes X_1^\pm \\ \Delta(X_2^+) &= X_2^+ \otimes (q^{\frac{1}{2}} s)^{-\frac{1}{2}H_1} (q^{\frac{3}{2}} s^{-1})^{\frac{1}{2}K} + (q^{\frac{1}{2}} s)^{\frac{1}{2}H_1} (q^{\frac{3}{2}} s^{-1})^{-\frac{1}{2}K} \otimes X_2^+ \\ \Delta(H_1) &= H_1 \otimes 1 + 1 \otimes H_1 \\ \Delta(K) &= K \otimes 1 + 1 \otimes K. \end{aligned} \quad (16)$$

The co-product of X_{12}^+ may be obtained from these relations, commutation relations, and the fact that Δ is a homomorphism of algebra.

The antipode: This is also inherited from the $U_q(\mathfrak{gl}(3))$ and our choice of parameters.

$$\begin{aligned} S(H_1) &= -H_1 & S(K) &= -K & S(X_1^-) &= -q^{-2} X_1^- \\ S(X_1^+) &= -q^2 X_1^+ & S(X_2^+) &= -q^2 X_2^+ & S(X_{12}^+) &= -q^4 X_{12}^+. \end{aligned} \quad (17)$$

At this point it is convenient to introduce another notation for the generators which reflects their “geometric” meaning better,

$$(K, J_3, J_+, J_-, P_1, P_2) := (K, H_1, X_1^+, X_1^-, X_{12}^+, X_2^+). \quad (18)$$

Now we generalize this method to higher dimensions. The procedure is as follows. We use the multi-parameteric deformation of $U(\mathfrak{gl}(n+1))$ which depends on $1 + \frac{1}{2}n(n+1)$ parameters

$$\{q, s_{ij}, i, j = 1, 2, \dots, n\} \quad (19)$$

and n^2 generators

$$\{H_i, I, X_{ij}^\pm, i \leq j = 1, 2, \dots, n\}. \quad (20)$$

Here, for notational convenience we have named $X_{ii}^\pm := X_i^\pm$, which are simple roots, and $X_{ij}^\pm$ for $i \neq j$ are composite roots. The matrices of the generators, in the un-deformed case are:

$$X_{ij}^+ = e_{ij+1}, \quad X_{ij}^- = e_{j+1i}, \quad H_i = e_{ii} - e_{i+1i+1}, \quad I = \sum_{i=1}^{n+1} e_{ii}. \quad (21)$$

In the above expressions, e_{ij} are the standard basis of matrices, with entries:

$$(e_{ij})_{mn} = \delta_{im} \delta_{jn}. \quad (22)$$

$U_q(\mathfrak{gl}(n))$ is generated by the following subset of generators:

$$\{H_i, K, X_{ij}^\pm, \quad i \leq j = 1, 2, \dots, n-1\} \quad (23)$$

where

$$K = \frac{1}{n+1} \sum_{i=1}^n i H_i + \frac{n}{n+1} I. \quad (24)$$

A basis for $\mathfrak{igl}(n)$ is obtained by adding to the set (23) the following translators:

$$P_i := X_{in}^+ \quad i = 1, 2, \dots, n. \quad (25)$$

In the quantum case also, we introduce this set as the generating set of the multi-parametric $U_q(\mathfrak{igl}(n))$.

$$\mathcal{B} = \{K, H_i, X_{ij}^\pm, \quad i \leq j = 1, 2, \dots, n-1, P_k, \quad k \leq n\} \quad (26)$$

$\mathcal{B}$ is a subset of the multi-parametric $U_q(\mathfrak{gl}(n+1))$. It is straightforward to check that this set is closed under “commutation”.

Looking at the co-product one sees that H_n and I appear in some of the terms, e.g. $\Delta(X_i^\pm)$. However, we have the freedom of fixing the parameters s_{ij} and s_i in such a way that only H_j for $j < n+1$ and K appear in these expressions. If we fix the parameters s_{ij} and s_i as below

$$s_i = q^{\frac{1}{2}\delta_{in}} \prod_{j=1}^n s_{ij}^{a_{jn}^{-1}} \quad (27)$$

the generators H_n and I disappear from the co-products. This will fix n parameters and we are left with $1 + \frac{1}{2}n(n-1)$ free parameters. We have checked that co-unity and antipode also respect this splitting of generators of $U_q(\mathfrak{gl}(n+1))$ for the above mentioned choice of parameters.

In summary, using the multi-parametric deformation of $U(\mathfrak{gl}(n+1))$, a special choice of the parameters lead to a Hopf algebra that contains a sub-Hopf-algebra generated by $\mathcal{B}$, which we call it $U_q(\mathfrak{gl}(n))$. It depends on $1 + \frac{1}{2}n(n-1)$ parameters. $U_q(\mathfrak{gl}(n))$ is generated by the set of generators of its subalgebra $U_q(\mathfrak{gl}(n-1))$ and n translations $P_i := X_{i,n}^+$. The commutation relations are the following, note that we have written the sub-algebra $\mathfrak{gl}(n)$ in the Chevalley basis

$$\begin{aligned} [H_i, H_j] &= 0 & [K, \dots] &= 0 \\ [H_i, X_j^\pm] &= \pm a_{ij} X_j^\pm & [X_i^+, X_j^-] &= \delta_{ij} [H_i] \end{aligned} \quad (28)$$

$$(X_{i\pm 1}^\pm)^2 X_i^\pm - (q + q^{-1}) X_{i\pm 1}^\pm X_i^\pm X_{i\pm 1}^\pm + X_i^\pm (X_{i\pm 1}^\pm)^2 = 0 \quad (29)$$

$$\begin{aligned} [H_i, P_j] &= \sum_{k=j}^n a_{ik} P_k & [K, P_i] &= P_i \\ [X_i^+, P_i]_q &= 0 & [P_i, X_j^-] &= \delta_{ij} q^{-\frac{1}{2}} P_{i+1} q^{H_i} \\ [P_i, X_{i-1}^+]_q &= P_{i-1} & [P_i, X_j^+] &= 0, \quad |i-j| \leq 2 \end{aligned} \quad (30)$$

The co-products are

$$\Delta(H_i) = H_i \otimes 1 + 1 \otimes H_i \quad (31)$$

$$\Delta(K) = K \otimes 1 + 1 \otimes K. \quad (32)$$

$$\begin{aligned} \Delta(X_i^\pm) &= X_i^\pm \otimes q^{\frac{1}{2}H_i} \left(\prod_{j=1}^n s_{ij}^{\pm \left[\sum_{k=1}^{n-1} \left(\frac{k(n-1)}{n} \right) - (k-1)\theta_{jk} \right] H_k + (1-j+\frac{1}{n})K} \right) + \\ & \quad q^{-\frac{1}{2}H_i} \left(\prod_{j=1}^n s_{ij}^{\mp \left[\sum_{k=1}^{n-1} \left(\frac{k(n-1)}{n} \right) - (k-1)\theta_{jk} \right] H_k + (1-j+\frac{1}{n})K} \right) \otimes X_i^\pm \end{aligned} \quad (33)$$

$$\begin{aligned} \Delta(P_n) &= P_n \otimes q^{-\frac{1}{2} \sum_{k=1}^{n-1} \frac{k}{n} H_n} \left(\prod_{j=1}^n s_{nj}^{\sum_{k=1}^{n-1} \left(\frac{k(n-1)}{n} \right) - (k-1)\theta_{jk}} \right)^{H_k} \left(q^{\frac{n+1}{2n}} \left(\prod_{j=1}^n s_{nj}^{1-j+\frac{1}{n}} \right) \right)^K + \\ & \quad q^{\frac{1}{2} \sum_{k=1}^{n-1} \frac{k}{n} H_n} \left(\prod_{j=1}^n s_{nj}^{\sum_{k=1}^{n-1} \left(\frac{-k(n-1)}{n} \right) + (k-1)\theta_{jk}} \right)^{H_k} \left(q^{-\frac{n+1}{2n}} \left(\prod_{j=1}^n s_{nj}^{-1+j-\frac{1}{n}} \right) \right)^K \otimes P_n \end{aligned} \quad (34)$$

for computing these co-products one must use (14). The co-products of the other P_i s can be obtained by using the fact that Δ is a homomorphism of the algebra and using the following relation

$$P_i = [\dots [P_n, X_{n-1}^+]_q, X_{n-2}^+]_q, X_{n-3}^+]_q, \dots X_i^+]_q. \quad (35)$$

After the submission of this paper we have been noted that multiparametric deformation of the algebra of functions $IGL_q(n)$ and its differential calculus have been studied in [15] where the authors found constraints on the parameters by taking dilatations in the center of the algebra.

References

- [1] Celeghini E., R. Giachetti, E. Sorace and M. Tarilini, J. Math. Phys. **32**, 1159 (1991)
- [2] Woronowicz, S.L., Commun. Math. Phys. **136**, 399 (1991)
- [3] Woronowicz, S.L., Lett. Math. Phys. **23**, 251 (1991)
- [4] Schliker M., W. Weich and R. Weixler, Z. Phys. C **53**, 79 (1992)
- [5] Schliker M., W. Weich and R. Weixler, Lett. Math. Phys. **27**, 217 (1993)
- [6] Castellani L., Phys. Lett. **B298**, 335 (1993); Castellani L., Lett. Math. Phys. **30**, 233 (1994);
Castellani L., Phys. Lett. **B327** (1994)
- [7] Aghamohammadi A., S. Rouhani and A. Shariati, To appear in J. Phys. A.
- [8] Manin Yu. I., Commun. Math. Phys. **123**, 163 (1989)
- [9] Reshetikhin N. Yu., Lett. Math. Phys. **20**, 331 (1990)
- [10] Saudbery A., To appear in J. Algebra
- [11] Schirrmacher A., J. Wess and B. Zumino, Z. Phys. C **49**, 317 (1991)
- [12] Schirrmacher A., Z. Phys. C **50**, 321 (1991)
- [13] Dobrev V. K. and P. Parashar, J. Phys. A **26**, 6991 (1993)
- [14] Karimipour V. and A. Aghamohammadi, J. Math. Phys. **34**, 2561 (1993)
- [15] Aschieri P. and L. Castellani, hep-th/9408031